



How to use your FASENRA Pen



Injecting with the FASENRA Pen is a simple 2-step process:

1. Prepare your supplies. 2. Inject.

Your healthcare provider will give you a demonstration before you begin, and can answer any questions you may have.

Prepare

Each single-use pen contains one dose of FASENRA. When you are ready to start, take the pen out of the refrigerator. Let it warm to room temperature—usually about 30 minutes. In the meantime, you can get the other things you will need.

Alcohol wipes



Cotton balls




Bandage



Sharps Disposal Container



Before you begin, wash your hands well with soap and water. **Choose your injection site**, and clean it in a circular motion with an alcohol wipe.



If you are giving yourself the injection, the **recommended injection site** is the front of your thigh or the lower part of your stomach (abdomen). A caregiver may inject you in the upper-arm, thigh, or stomach. Do not try to inject yourself in the arm.

For each injection, choose a different site that is at least 1 inch away from your last injection.

■ Self-admin and caregiver injection site

▨ Caregiver-only injection site

Do not inject into the 2-inch area around your belly button, or where the skin is tender, bruised, scaly, or hard, into scars or damaged skin, or through clothing.

Please see additional Important Safety Information and full [Prescribing Information](#), including [Patient Information](#) and [Instructions for Use](#).

Inject

Before you inject, please review the steps below.



1. Remove the cap. Be careful not to touch the needle or push on the green needle guard.



2. Line the FASENRA Pen up at a 90-degree angle with the injection site. You may gently pinch the skin surrounding your injection site, but this is not a requirement.



3. Press down firmly and hold. You will hear a click signaling that the injection has begun. The green plunger will start to fill the viewing window. Hold the FASENRA Pen in place for 15 seconds and do not change the position of the pen.

You will hear a second click that signals the injection has finished. The green plunger should fill the viewing window.



4. Lift the FASENRA Pen straight up. The needle guard will slide down and lock into place over the needle. You may see a small amount of blood or liquid at the injection site. Gently apply pressure to the area with a cotton ball until the bleeding stops. Do not rub the injection site. If needed cover it with a bandage.

To dispose of the used pen safely, put it in your Sharps Disposal Container right after use.

For additional resources and helpful information about the FASENRA Pen and self-injection, visit fasenrapen.com, or call 1-833-360-HELP (4357).

Please see additional Important Safety Information and full [Prescribing Information](#), including [Patient Information](#) and [Instructions for Use](#).

IMPORTANT SAFETY INFORMATION

Do not use FASENRA if you are allergic to benralizumab or any of the ingredients in FASENRA.

Do not use to treat sudden breathing problems.

FASENRA may cause serious side effects, including:

- **allergic (hypersensitivity) reactions, including anaphylaxis.** Serious allergic reactions can happen after you get your FASENRA injection. Allergic reactions can sometimes happen hours or days after you get your injection. Tell your healthcare provider or get emergency help right away if you have any of the following symptoms of an allergic reaction:
 - swelling of your face, mouth and tongue
 - breathing problems
 - fainting, dizziness, feeling lightheaded (low blood pressure)
 - rash
 - hives

Before using FASENRA, tell your healthcare provider about all of your medical conditions, including if you:

- are taking oral or inhaled corticosteroid medicines. **Do not** stop taking your corticosteroid medicines unless instructed by your healthcare provider. This may cause other symptoms that were controlled by the corticosteroid medicine to come back.
- have a parasitic (helminth) infection.
- are pregnant or plan to become pregnant. It is not known if FASENRA will harm your unborn baby. Tell your healthcare provider if you become pregnant during your treatment with FASENRA.
 - There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to FASENRA during pregnancy. Healthcare providers can enroll patients or encourage patients to enroll themselves by calling 1-877-311-8972 or visiting www.mothers-to-baby.org/fasenra.
- are breastfeeding or plan to breastfeed. It is not known if FASENRA passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you use FASENRA.
- are taking prescription and over-the-counter medicines, vitamins, or herbal supplements.

Do not stop taking your other asthma medicines unless instructed to do so by your healthcare provider.

The most common side effects of FASENRA include: headache and sore throat. These are not all the possible side effects of FASENRA.

APPROVED USE

FASENRA is a prescription medicine used with other asthma medicines for the maintenance treatment of asthma in people 12 years and older whose asthma is not controlled with their current asthma medicines. FASENRA helps prevent severe asthma attacks (exacerbations) and may improve your breathing. Medicines such as FASENRA reduce blood eosinophils. Eosinophils are a type of white blood cell that may contribute to your asthma.

FASENRA is not used to treat other problems caused by eosinophils and is not used to treat sudden breathing problems. Tell your health care provider if your asthma does not get better or if it gets worse after you start treatment with FASENRA.

It is not known if FASENRA is safe and effective in children under 12 years of age.

You may report side effects related to AstraZeneca products by clicking [here](#).

Please see accompanying full [Prescribing Information](#), including [Patient Information](#) and [Instructions for Use](#).